

Methyl jasmonate-dependent senescence of cotyledons in *Ipomoea nil*

E. Wilmowicz^{1,2} · A. Kućko^{1,2} · K. Frankowski¹ · M. Świdziński^{2,3} · K. Marciniak¹ · J. Kopcewicz¹

Received: 12 January 2016/Revised: 9 August 2016/Accepted: 22 August 2016/Published online: 26 August 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract Jasmonic acid methyl ester (JAME) has been recently shown to play a crucial role in many physiological processes. In this paper, we focused on cotyledon senescence in *Ipomoea nil* and revealed that JAME and darkness are the main factors stimulating the process examined. What is more, we showed that mefenamic acid (a jasmonate biosynthesis inhibitor) reverses the stimulatory effect of darkness on senescence. In plants growing under dark conditions, stimulation of *JASMONIC ACID CARBOXYL METHYLTRANSFERASE* (*InJMT*) expression and, consequently, an increase in JAME content, have been observed. In turn, the level of jasmonic acid (JA) gradually decreased. Moreover, dark-grown seedlings demonstrated a lower PSII functional activity and a reduced chlorophyll content and autofluorescence. All of these data suggest that JAME is a signal molecule controlling the senescence of cotyledons in *I. nil*.

Keywords *Ipomoea nil* · Jasmonates · Jasmonic acid methyl ester · Senescence · Jasmonic acid carboxyl methyltransferase

Introduction

The senescence of cotyledons is the final stage of their development. This process is an integrated response of these organs to age information and other environmental signals, both internal and external, such as stresses of salinity, nutrient limitation or darkness (Bouchard and Yamasaki 2008; Corpas et al. 2009; Du et al. 2014; Ma et al. 2008; Neill et al. 2008; Xuan et al. 2010; Zhao et al. 2009). Although there is no single universal mechanism for regulating senescence, some characteristic stages of that process can be distinguished. Initially, thylakoid membrane structures are damaged, the photosynthetic rate drops as a result of chlorophyll degradation, and the catabolic process stops (Pružinská et al. 2005; Troncoso-Poncea et al. 2013; Wrisher et al. 2009; Zhang et al. 2010). Subsequently, the ER, ribosomes, proteins, lipids and nucleic acids are decomposed. In the final stage of cotyledon senescence, the mitochondria are degraded, oxidative phosphorylation intensity is decreased, and the tonoplast and nuclear membrane are interrupted, all of which lead to irreversible changes in the structures of the cytoplasm and the nucleus, as well as to the accumulation of secondary metabolites (Lim et al. 2007; Peterman and Siedow 1985). Although cotyledon senescence has been studied extensively (Ananieva et al. 2008a, b; Kanazawa et al. 2000; Krul 1974; McKersie et al. 1987; Peterman and Siedow 1985; Rukes and Mulkey 1993; Watanabe et al. 1994), its underlying regulatory network is unclear in comparison to leaf senescence (Lim et al. 2007). Numerous studies have

Communicated by P. K. Nagar.

Electronic supplementary material The online version of this article (doi:10.1007/s11738-016-2244-1) contains supplementary material, which is available to authorized users.

✉ E. Wilmowicz
emwil@umk.pl

- ¹ Chair of Plant Physiology and Biotechnology, Nicolaus Copernicus University, 1 Lwowska Street, 87-100 Toruń, Poland
- ² Centre for Modern Interdisciplinary Technologies, Nicolaus Copernicus University, 4 Wileńska Street, 87-100 Toruń, Poland
- ³ Department of Cell Biology, Nicolaus Copernicus University, 1 Lwowska Street, 87-100 Toruń, Poland

shown that leaf senescence correlates with a progressive decline in the photosynthetic rate, which is accompanied by Rubisco and chlorophyll *a/b* binding protein (CAB) degradation (Avice and Etienne 2014; Gombert et al. 2006; Lu et al. 2002; Van der Graaff et al. 2006). There are crucial photosynthetic parameters commonly known as senescence physiological markers e.g., chlorophyll content, rate of photosynthesis (Gan 2004). In addition, the transcriptional activity of senescence-associated (SAGs) genes increased in leaves before or during senescence in different species (Breeze et al. 2011; Buchanan-Wollaston et al. 2005). Many SAGs are similar to genes probably involved in the breakdown and mobilization of nutrients, such as proteases, RNases, or glutamine synthetases (Lin et al. 2015). An essential role in coordinating anatomical, physiological and biochemical changes accompanying senescence is attributed to plant hormones, such as cytokinin, ethylene, abscisic acid, auxin and jasmonates (JAs) (Ananieva et al. 2008a; Ellis et al. 2005; Hung and Kao 2004; Jing et al. 2008; Okushima et al. 2005). Hormonal pathways are involved in cotyledon senescence—process that may be age-controlled or stimulated by plant responses to stresses. Such hormones may also affect the synthesis and signaling pathways of other to eventually trigger the expression of stress-responsive genes, which in turn influences cotyledon senescence (Lim et al. 2007).

Lipid-derived JAs arise from the enzymatic oxygenation of 18- and 16-carbon triunsaturated fatty acids. The biosynthesis of these hormones encompasses three cellular compartments: chloroplast, peroxisome and cytosol. The first stage occurs in the chloroplasts, where, thanks to the activity of 13-LOX (lipoxygenase), AOS (allene oxide synthase) and AOC (allene oxide cyclase), the 18:3 or 16:3 acid is oxidized and OPDA (12-oxophytodienoic acid) or dnOPDA (dinor-OPDA) is formed, respectively. Both compounds are transported into peroxisomes and with the participation of OPR3 (12-oxophytodienoate reductase), ACX (acyl-CoA oxidase), MFP (multifunctional protein) and KAT (L-3-ketoacyl-CoA thiolase) are first reduced, and then β -oxidized, which ends in the formation of the (+)-7-iso-jasmonic acid (JA) in the cytosol (Wasternack and Kombrink 2010). Then, JA can be metabolized by esterification into methyl (+)-7-iso-jasmonate (JAMe) or by amino acid conjugation to form (+)-7-iso-jasmonoyl-isoleucine (JA-Ile). The second is considered as the active form of the hormone. In addition, JA can be hydroxylated, sulfated, as well as glucosylated (Acosta and Farmer 2010).

The use of molecular biology techniques in research on the JA signal transduction pathway and regulation of its metabolism has led to the discovery that JA derivatives can be also biologically active in many physiological processes. Studies performed in many plant species show that JAMe, formed in the reaction catalyzed by JASMONIC

ACID CARBOXYL METHYLTRANSFERASE (JMT), regulates the sprouting of seeds, participates in proper flower morphogenesis and flower induction, and is involved in responses to stress, which also includes promoting effect on leaf senescence (Ananieva et al. 2007; Kęsy et al. 2011; Maciejewska and Kopcewicz 2002; Maciejewska et al. 2004; Norastehnia et al. 2007; Wang 1998; Wilson et al. 2011).

In our former investigation, we showed that JAMe is an inhibitor of photoperiodic flower induction in model short-day plant (SDP) *Ipomoea nil*, in which the cotyledons are responsible for the perception of light stimuli (Kęsy et al. 2011; Maciejewska and Kopcewicz 2002; Maciejewska et al. 2004). Bearing in mind the involvement of JAMe in the control of many processes, we conducted physiological experiments on the efficiency of PSII, changes in the jasmonate level, chlorophyll content and autofluorescence. What is more, we performed identification and expression analyses of the *InJMT* gene participating in JA metabolism accompanying cotyledon senescence.

Materials and methods

Plant material

We prepared plant material (*I. nil*, Chois cv. Violet; Marutane Seed Co., Kyoto, Japan) according to Frankowski et al. (2009). The seedlings were grown in phytotrons under the conditions described by Wilmowicz et al. (2008). On 6th day of cultivation, half of the seedlings were exposed to darkness (D), whereas the remainder to continuous light (CL). The plants were watered with identical amounts of tap water every 2 days (Supplementary Fig. 1).

Plants growing in darkness were treated with mefenamic acid (MEF), a jasmonate biosynthesis inhibitor, at a concentration of 100 μ M in 0.05 % Tween 20 (v/v), whereas plants cultivated under CL were treated with JA or JAMe at a concentration of 100 μ M in 0.05 % Tween 20 (v/v). The control plants were treated with 0.05 % Tween 20 (v/v). All solutions were applied by small, soft brushes to the cotyledons (about 50 μ L per plant). All treatments were repeated every day at the same time starting from 6th day of cultivation for eight subsequent days. Before each treatment, the cotyledons from darkness and those after MEF application were collected (day 1, 2, 4, 6 and 8). In addition, to monitor the ongoing process of aging, the amount of quantum efficiency of PSII in the cotyledons was simultaneously investigated. The collected material was placed immediately in frozen liquid nitrogen and stored at -80°C . Each experiment was repeated at least

three times (biological repetitions). All data are presented as mean \pm standard error (SE).

Determination of chlorophyll content and fluorescence measurement

The chlorophyll content was determined according to Glazińska et al. (2014) with the method using Opti-Sciences CCM-200 (Opti-Sciences, Inc., USA).

Chlorophyll fluorescence changes are a useful index to measure photosynthetic efficiency. The fluorescence parameters were F_0 and F_m —the initial or variable fluorescence yield, respectively, while the F_v/F_m ratio indicated a maximum potential of the plant's photosynthetic ability. In our investigation, we used chlorophyll fluorometer OS-30P (Opti-Sciences, Inc., USA) according to Weng (2006). The data obtained were subjected to a statistical analysis and presented as mean \pm SE. Student's t test was used to calculate the significant differences as compared with the control.

Chloroplast detection using confocal microscopy

The collected tissue fragments were analyzed with using Nikon Eclipse TE300 confocal laser scanning inverted microscope. Obtained results were registered with an argon ion laser and He–neon laser emitting light with a wavelength of 488 nm (blue excitation and green fluorescence) and 543 nm (green excitation and red fluorescence), respectively. A mid pinhole, a long exposure time (90 s) and 100 \times (numerical aperture, 1.4) Plan Apochromat DCI H oil immersion lens were used in the analyses. We collected pairs of images in the green and red channels simultaneously. The EZ 2000 Viewer software package (Nikon Europe BV, Badhoevedorp, the Netherlands) was used for results documentation and analysis.

Determination of endogenous JAs

Endogenous jasmonates (jasmonic acid—JA, methyl jasmonate—JAMe) were determined with the method described by Wilmowicz et al. (2014) with modifications. Internal standards were 100 ng d₂-JAMe and 100 ng d₅-JA added to the crude extract. GC/MS-selected ion monitoring was performed by monitoring m/z 193, 195, 198, 224, 226 and 229.

Molecular cloning of *InJMT* cDNA

A full sequence of *InJMT* was obtained in PCR reactions using specific primers 5'-TTCCCCGAAAGCATGGGCA-3' (forward); 5'-TCGGTTGTAGGATCGCCGGA-3' (reverse) designed for EST fragments of *InJMT* (identified EST

sequences of *P. nil* in the NCBI database: no. BJ560292.1 and CJ756546.1) (Supplementary Fig. 2). The reaction mixture was subjected to the following PCR conditions: 95 °C for 5 min, 1 cycle; 95 °C for 1 min, 62–50 °C for 30 s, 72 °C for 45 s for 15 cycles, followed by 1 cycle of incubation at 72 °C for 7 min. An amplified cDNA fragment was isolated from an agarose gel with the GeneMATRIX Agarose-Out DNA Purification Kit (EurX, Gdańsk, Poland) and sequenced by “Genomed S.A.” (Warsaw, Poland). The full-length *InJMT* sequence has been deposited in the GenBank database.

The bioinformatical analysis and phylogenetic tree calculation of identified cDNA was made with using ClustalW (<http://www.ebi.ac.uk/clustalw>), BLAST 2.2.25 (<http://www.ncbi.nlm.nih.gov/BLAST>) and ExPASy (<http://www.expasy.org>) and the CLC Main Workbench v.5.6.1 program.

Quantitative real-time PCR (qPCR)

The gene expression analysis was performed by Real-Time PCR (RT-qPCR) with a LightCycler 2.0 Carousel-Based System and LightCycler TaqMan Master Kit (ROCHE Diagnostics GmbH, Germany). The cDNA templates were prepared according to Frankowski et al. (2015), whereas gene-specific primers and UPL probes for the *InJMT* (GenBank acc. no. KF573520.1) were designed using Universal ProbeLibrary Assay Design Center (<http://www.roche-applied-science.com/sis/rtpcr/upl>) (Supplementary Table 1). *InACT4* (GenBank acc. no. HM802138.2) was used as a reference gene. qPCR reaction was performed according to Frankowski et al. (2015). Three independent replicate RNA preparations were used for the analyses, and data are shown as mean \pm SE of three samples for each stage.

Results

Darkness and JAMe induce cotyledon senescence in *I. nil*

The progress of senescence was monitored by observing the inhibition of the functional activity of PSII (F_v/F_m), loss of chlorophyll and reduction of its autofluorescence. Our results showed that darkness promotes senescence through chlorophyll degradation (Fig. 1b) and autofluorescence (Fig. 2b), as well as a gradual decrease of the efficiency of PSII (Fig. 1a). The effect observed is accompanied by the progressive senescence processes. The F_v/F_m value in dark-grown plants and in light-grown plants treated with JAMe is 15 and 20 % lower than in control plants, respectively (Fig. 1a). The same solution applied to the cotyledons of plants grown under light conditions




Fig. 1 Changes in the maximum quantum efficiency of PSII [variable fluorescence (F_v)/maximum fluorescence (F_m) ratio] (a) and chlorophyll content (b) in the cotyledons of *Ipomoea nil*. CL continuous light, D darkness, JA jasmonic acid, JAME jasmonic acid methyl ester, MEF mefenamic acid (a jasmonate biosynthesis inhibitor). Vertical bars indicate SE. Significant differences to the plants growing under light condition are indicated as ** $P < 0.01$ and * $P < 0.05$

decreased the autofluorescence of the chlorophyll (Fig. 2c). JA has no influence on senescence in light-grown plants (Fig. 1a, b). Nevertheless, the application of MEF (a jasmonate biosynthesis inhibitor) on the cotyledons of *I. nil* reverses the stimulatory effect of the darkness on the process discussed. These plants contain a higher chlorophyll concentration than the cotyledons of plants grown in darkness and have a higher F_v/F_m ratio value (Fig. 1a, b), and also, the chlorophyll contained in them demonstrates autofluorescence similar to that of the control light-grown plants (Fig. 2d).

Isolation of the *InJMT* cDNA

The full-length *InJMT* cDNA (GB acc. no KF573520.1) obtained using the RT-PCR technique is composed of 1403 bp and encodes 380 amino acids (Fig. 3a, b). The predicted protein sequence of InJMT contains all the

conserved domains characteristic of JMT identified in other plant species. A phylogenetic analysis of the amino acid sequence revealed that the predicted InJMT is very closely related to the JMT from *Solanum tuberosum* (StJMT, GB acc. no XP006341965.1) (Fig. 3c).


Senescence of *I. nil* cotyledons is accompanied by increase in the JAME content and *InJMT* transcriptional activity

To investigate the possible role of JAs in cotyledon senescence in *I. nil*, we analyzed variations in the endogenous JA and JAME level and the expression of *InJMT* coding for carboxyl methyltransferase during various stages of aging. The endogenous content of JAME in the cotyledons of *I. nil* growing under darkness is initially hardly any lower than that of JA (Fig. 4b). However, the subsequent days see a gradual growth in the JAME content in the cotyledons and a drop in the JA level. On day 8, the JAME content is over four times that of JA, and over twice that of JAME content in the control plants (Fig. 4b). The increased endogenous JAME level is correlated with a stimulation of the *InJMT* transcriptional activity (Fig. 4a). Moreover, plants growing in darkness and treated with the jasmonate biosynthesis inhibitor (MEF) showed decrease in the *InJMT* expression and JAME content (Fig. 4a, b).

Discussion

Cotyledons are organs forming during embryogenesis as initially heterotrophic, after which they become photosynthetically active, and then, their function is to provide nutrients for seedlings. The senescence of cotyledons, like in the case of other organs, is an integral part of their development; similar to leaf senescence, it is genetically programmed and occurs when they already fulfilled their role. However, this process can be stimulated prematurely by stress factors, both biotic, e.g., pathogen infection, and abiotic, e.g., extreme temperatures, drought, nutrient deficiency, and shading (Krul 1974; Quirino et al. 2000). An important role in regulating this process is played by phytohormones, which are signal molecules and/or coordinators in the transformations accompanying senescence, such as nutrient recycling, the dismantling of chloroplasts, the cell death as a consequence of DNA, RNA and proteins degradation (Krul 1974; Peterman and Siedow 1985; Quirino et al. 2000; Wada et al. 2009). Among these phytohormones, jasmonates are particularly noteworthy. These substances cause chlorophyll loss, degradation of Rubisco and inhibition of its biosynthesis. They can also induce the disorganization of the thylakoid membrane system in chloroplasts, affect the functional activity of

Fig. 2 Visualization of chlorophyll autofluorescence in the cotyledons of *Ipomoea nil* growing under: continuous light (a), dark conditions (b), treated with jasmonic acid methyl ester after transfer to continuous light (c), treated with mefenamic acid (inhibitor of jasmonate biosynthesis) after transfer to darkness (d). Bars 200 μ m



PSII, decrease the rates of transpiration and photosynthesis, as well as cell up-regulates several SAGs shown in different plants (Ananieva et al. 2004b; Beltrano et al. 1998; He and Gan 2001; He et al. 2002; Liu et al. 2015; Weaver et al. 1998; Weidhase et al. 1987a, b; Woo et al. 2001; Oh et al. 1996). Jasmonates have been widely shown to be powerful promoters of leaf senescence (Chen and Kao 1998; Chou and Kao 1992; Hung and Kao 1996; Tsai et al. 1996; Ueda and Kato 1981; Weidhase et al. 1987a), but their participation in cotyledon senescence is still poorly understood (Ananieva et al. 2004a, 2007). In the present study, we evaluate the effects of darkness and JAs on the senescence of *I. nil* cotyledons.


We showed that darkness causes progressive reduction in the content of chlorophyll in the cotyledons (Fig. 1b) and decreases its autofluorescence (Fig. 2c), diminishing the F_v/F_m ratio, as well (Fig. 1a). In many plant species, the stimulating effect of darkness on senescence processes was observed (Biswal et al. 1983; Biswal and Biswal 1984; Oh et al. 1996; Weaver et al. 1998). Reducing the endogenous jasmonate content by applying the jasmonate biosynthesis inhibitor reverses dark-stimulated cotyledon senescence in *I. nil* (Figs. 1a, b, 2d), which confirms that these phytohormones play a key role in regulating this process. Although the conjugate of JA and the amino acid isoleucine (Ile) has been reported to be a bioactive form of jasmonates, the results of substantial physiological and

molecular research do not eliminate a specific role played by other jasmonates, such as JAMe, in controlling of growth and developmental processes (Noir et al. 2013; Pak et al. 2009; Staswick 1992). Moreover, JAMe is a plant volatile regulator and is thought to be a potential candidate for intra- and intercellular signal transducers mediating jasmonate-dependent plant responses because of its ability to diffuse through the membranes (Seo et al. 2001; Cheong and Choi 2003).

JAMe applied to the cotyledons of light-grown *I. nil* accelerates their senescence. Both the chlorophyll content (Fig. 1b) and the F_v/F_m ratio (Fig. 1a) assume values lower than in plants with dark-stimulated senescence. No such effect was observed following JA applications (Fig. 1a, b). JAs, i.e., both JA and JAMe, are promoters of leaf senescence upon their exogenous application (Creelman and Mullet 1997; Wasternack and Hause 2002). JAMe is the most effective stimulator of leaf senescence in the leaves of wild-type and mutant *Arabidopsis* (Oh et al. 1996; Weaver et al. 1998; Woo et al. 2001), leaf segments of barley (Reinbothe et al. 1993, 1997; Weidhase et al. 1987a) and intact cotyledons of *Cucurbita pepo* (zucchini) (Ananieva et al. 2004b).

Darkness leads to increased concentrations of endogenous JAMe and decreases the pool of free JA (Fig. 4b), which shows that JAMe is a cotyledon senescence stimulator in *I. nil*. If a change in the JAMe level is important for

Fig. 3 Identification of full-length *InJMT* cDNA. Products of PCR with specific primers (**a**). The coding sequence of *InJMT* and the deduced amino acid sequence (**b**). Subsequent nucleotide positions are marked on the *left side* of the figure, and amino acid positions on the *right*. The translation initiation point (the start codon) and termination point (the stop codon) are in **boldface** type. *Small letters* are used for regions not subject to translation. Domains responsible for methyl, salicylate, SAM (*S*-adenosyl methionine) and SAH (*S*-adenosyl-L-homocysteine) group binding are marked with *light gray*, *dark gray*, or are *underlined*, respectively. A phylogenetic analysis of *InJMT* with selected plant *JMT* sequences (**c**). Accession numbers of the sequences used to build the tree were as follows: RcJMT (XP002510424.1), SIJMT (XP004238275.1), StJMT (XP006341965.1), InJMT (AHA06001.1), VvJMT (XP002281588.1), TcJMT (XP007017600.1), PtJMT (XP002307671.1)



explaining cotyledon senescence, it is crucial to study the expression level of the *InJMT* gene that encodes a cytosolic protein involved in the methylation of JA. In the present investigation, we identified this gene in *I. nil* (Fig. 3).

Based on a comparison between the deduced amino acid sequences of *InJMT* and *JMTs* from other plant species, it was found that *InJMT* contains all of the characteristic motifs present in SAM and SAM-like methyltransferases,




Fig. 4 The expression level of *InJMT* (related to *InACT4*) in the cotyledons of *I. nil* seedlings during darkness conditions and following MEF treatment (**a**). Dark-gray bars D conditions, gray bars MEF applied during darkness conditions. The endogenous jasmonates (JA jasmonic acid, JAMe jasmonic acid methyl ester) level in the cotyledons growing under darkness conditions and JAMe after MEF treatment (**b**). Mean \pm SE $n = 3$. The thick horizontal line under the time axis indicates duration of the dark period. For JA, significant differences to the plants growing under SD condition are indicated as ^{bb} $P < 0.01$ and ^b $P < 0.05$. For JAMe, significant differences to the plants growing under SD condition are indicated as ^{**} $P < 0.01$ and ^{*} $P < 0.05$. For JAMe, after MEF treatment, significant differences to the plants growing under darkness condition are indicated as ^{aa} $P < 0.01$ and ^a $P < 0.05$

SAM/SAH binding residues, salicylate/JA binding residues and additional active site residues (Fig. 3b, c), which are essential for its activity (Martin and McMillan 2002). Bearing this in mind, it can be suspected that the gene encodes a functional enzymatic protein. A progressing senescence of *I. nil* cotyledons is correlated with a growth in *InJMT* expression (Fig. 4a), which elevates the JAMe content (Fig. 4b). The drop in the endogenous JAMe content on day 8 of darkness may be caused by the exhaustion of the available substrate for the enzymatic activity of *InJMT*.

These results suggest that the accelerated cotyledon senescence in dark-grown *I. nil* is correlated with an

increased *InJMT* expression level and, consequently, a growth in the concentration of JAMe which plays the role of a signal molecule in regulating this process.

Author contribution statement Emilia Wilmowicz and Agata Kućko designed and carried out all the experiments and wrote the manuscript. Kamil Frankowski helped with gene identification and qPCR reactions, Michał Świdziński conducted the microscopy experiments, Katarzyna Marciniak measured the chlorophyll content and quantum efficiency of PS II, Jan Kopcewicz and Agata Kućko were responsible for preparing the final version of manuscript. All authors were responsible for analysis and interpretation of obtained results.

Acknowledgments This project was supported by Nicolaus Copernicus University Grant No. 1233-B and funds provided by Nicolaus Copernicus University (Toruń, Poland) for the research program of the Chair of Plant Physiology and Biotechnology.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Acosta IF, Farmer EE (2010) Jasmonates. Arabidopsis Book 8:e0129
- Ananieva K, Malbeck J, Kamínek M, Van Staden J (2004a) Methyl jasmonate down-regulates endogenous cytokinin levels in cotyledons of *Cucurbita pepo* (zucchini) seedlings. *Physiol Plant* 122:496–503
- Ananieva K, Malbeck J, Kamínek M, Van Staden J (2004b) Changes in endogenous cytokinin levels in cotyledons of *Cucurbita pepo* (zucchini) during natural and dark induced senescence. *Physiol Plant* 122:133–142
- Ananieva K, Ananieva ED, Mishev K, Georgieva K, Malbeck J, Kamínek M, Van Staden J (2007) Methyl jasmonate is a more effective senescence-promoting factor in *Cucurbita pepo* (zucchini) cotyledons when compared with darkness at the early stage of senescence. *J Plant Physiol* 164:1179–1187
- Ananieva K, Ananieva ED, Doncheva S, Georgieva K, Tzvetkova N, Kamínek M, Motyka V, Dobrev P, Gajdosová S, Malbeck J (2008a) Senescence progression in a single darkened cotyledon depends on the light status of the other cotyledon in *Cucurbita pepo* (zucchini) seedlings: potential involvement of cytokinins and cytokinin oxidase/dehydrogenase activity. *Physiol Plant* 134:609–623
- Ananieva K, Ananieva ED, Mishev K, Georgieva K, Tzvetkova N, Van Staden J (2008b) Changes in photosynthetic capacity and polypeptide patterns during natural senescence and rejuvenation of *Cucurbita pepo* L. (zucchini) cotyledons. *Plant Growth Regul* 54:23–29
- Avicé JC, Etienne P (2014) Leaf senescence and nitrogen remobilization efficiency in oilseed rape (*Brassica napus* L.). *J Exp Bot* 65:3813–3824
- Beltrano J, Ronco MG, Montaldi ER, Carbone A (1998) Senescence of flag leaves and ears of wheat hastened by methyl jasmonate. *J Plant Growth Regul* 17:53–57

- Biswal UC, Biswal B (1984) Photocontrol of leaf senescence. *Photochem Photobiol* 39:875–879
- Biswal B, Choudhury NK, Sahu P, Biswal UC (1983) Senescence of detached fern leaves. *Plant Cell Physiol* 24:1203–1208
- Bouchard JN, Yamasaki H (2008) Heat stress stimulates nitric oxide production in *Symbiodinium microadriaticum*: a possible linkage between nitric oxide and the coral bleaching phenomenon. *Plant Cell Physiol* 49:641–652
- Breeze E, Harrison E, McHattie S, Hughes L, Hickman R, Hill C, Kiddle S, Kim YS, Penfold CA, Jenkins D, Zhang C, Morris K, Jenner C, Jackson S, Thomas B, Tabrett A, Legaie R, Moore JD, Wild DL, Ott S, Rand D, Beynon J, Denby K, Mead A, Buchanan-Wollaston V (2011) High-resolution temporal profiling of transcripts during *Arabidopsis* leaf senescence reveals a distinct chronology of processes and regulation. *Plant Cell* 23:873–894
- Buchanan-Wollaston V, Page T, Harrison E, Breeze E, Lim PO, Nam HG, Lin JF, Wu SH, Swidzinski J, Ishizaki K, Leaver CJ (2005) Comparative transcriptome analysis reveals significant differences in gene expression and signalling pathways between developmental and dark/starvation-induced senescence in *Arabidopsis*. *Plant J* 42:567–585
- Chen SJ, Kao CH (1998) Methyl jasmonate, ammonium, and leaf senescence in rice. *J Plant Physiol* 152:353–357
- Cheong JJ, Choi YD (2003) Methyl jasmonate as a vital substance in plants. *Trends Genet* 19:409–413
- Chou CM, Kao CH (1992) Methyl jasmonate, calcium, and leaf senescence in rice. *Plant Physiol* 99:1693–1694
- Corpas FJ, Hayashi M, Mano S, Nishimura M, Barroso JB (2009) Peroxisomes are required for in vivo nitric oxide accumulation in the cytosol following salinity stress of *Arabidopsis* plants. *Plant Physiol* 151:2083–2094
- Creelman RA, Mullet JE (1997) Biosynthesis and action of jasmonates in plants. *Annu Rev Plant Physiol Plant Mol Biol* 48:355–381
- Du J, Li M, Kong D, Wang L, Lv Q, Wang J, Bao F, Gong Q, Xia J, He Y (2014) Nitric oxide induces cotyledon senescence involving co-operation of the NES1/MAD1 and EIN2-associated ORE1 signalling pathways in *Arabidopsis*. *J Exp Bot* 65:4051–4063
- Ellis CM, Nagpal P, Young JC, Hagen G, Guilfoyle TJ, Reed JW (2005) AUXIN RESPONSE FACTOR1 and AUXIN RESPONSE FACTOR2 regulate senescence and floral organ abscission in *Arabidopsis thaliana*. *Development* 132:4563–4574
- Frankowski K, Kęsy J, Wojciechowski W, Kopcewicz J (2009) Light- and IAA-regulated ACC synthase gene (*PnACS*) from *Pharbitis nil* and its possible role in IAA-mediated flower inhibition. *J Plant Physiol* 166:192–202
- Frankowski K, Wilmowicz E, Kućko A, Zienkiewicz A, Zienkiewicz K, Kopcewicz J (2015) Profiling the *BLADE-ON-PETIOLE* gene expression in the abscission zone of generative organs in *Lupinus luteus*. *Acta Physiol Plant* 37:220. doi:10.1007/s11738-015-1972-y
- Gan S (2004) The hormonal regulation of senescence. In: Davies PJ (ed) *Plant hormones: biosynthesis, signal transduction and action*. Kluwer Academic Publishers, Dordrecht, pp 561–581
- Glazińska P, Wilmowicz E, Wojciechowski W, Frankowski K, Kopcewicz J (2014) Impact of *InMIR319* and light on the expression of *InTCP4* gene involved in the development of *Ipomoea nil* plants. *Acta Physiol Plant* 36:29–43
- Gombert J, Etienne P, Ourry A, Le Dily F (2006) The expression patterns of *SAG12/Cab* genes reveal the spatial and temporal progression of leaf senescence in *Brassica napus* L. with sensitivity to the environment. *J Exp Bot* 57:1949–1956
- He Y, Gan S (2001) Identical promoter elements are involved in regulation of the *OPR1* gene by senescence and jasmonic acid in *Arabidopsis*. *Plant Mol Biol* 47:595–603
- He Y, Fukushige H, Hildebrand DF, Gan S (2002) Evidence supporting a role of jasmonic acid in *Arabidopsis* leaf senescence. *Plant Physiol* 128:876–884
- Hung KT, Kao CH (1996) Promotive effect of jasmonates on the senescence of detached maize leaves. *Plant Growth Regul* 19:77–83
- Hung KT, Kao CH (2004) Hydrogen peroxide is necessary for abscisic acid-induced senescence of rice leaves. *J Plant Physiol* 161:1347–1357
- Jing HC, Schippers JH, Hille J, Dijkwel PP (2008) Ethylene induced leaf senescence depends on age-related changes and *OLD* genes in *Arabidopsis*. *J Exp Bot* 56:2915–2923
- Kanazawa S, Sano S, Koshiba T, Ushimaru T (2000) Changes in antioxidative enzymes in cucumber cotyledons during natural senescence: comparison with those during dark-induced senescence. *Physiol Plant* 109:211–216
- Kęsy J, Wilmowicz E, Maciejewska B, Frankowski K, Glazińska P, Kopcewicz J (2011) Independent effects of jasmonates and ethylene on inhibition of *Pharbitis nil* flowering. *Acta Physiol Plant* 33:1211–1216
- Krul WR (1974) Nucleic acid and protein metabolism of senescing and regenerating soybean cotyledons. *Plant Physiol* 54:36–40
- Lim PO, Kim HJ, Nam HG (2007) Leaf senescence. *Annu Rev Plant Biol* 58:115–136
- Lin M, Pang C, Fan S, Song M, Wei H, Yu S (2015) Global analysis of the *Gossypium hirsutum* L. Transcriptome during leaf senescence by RNA-Seq. *BMC Plant Biol* 12(15):43. doi:10.1186/s12870-015-0433-5
- Liu L, Li H, Zeng H, Cai Q, Zhou X, Yin Ch (2015) Exogenous jasmonic acid and cytokinin antagonistically regulate rice flag leaf senescence by mediating chlorophyll degradation, membrane deterioration, and senescence-associated genes expression. *J Plant Growth Regul*. doi:10.1007/s00344-015-9539-0
- Lu C, Koroleva OA, Farrar JF, Gallagher J, Pollock CJ, Deri TA (2002) Rubisco small subunit, chlorophyll *a/b*-binding protein and sucrose:fructan-6-fructosyl transferase gene expression and sugar status in single barley leaf cells in situ. Cell type specificity and induction by light. *Plant Physiol* 130:1335–1348
- Ma W, Smigel A, Tsai YC, Braam J, Berkowitz GA (2008) Innate immunity signaling: cytosolic Ca^{2+} elevation is linked to downstream nitric oxide generation through the action of calmodulin or a calmodulin-like protein. *Plant Physiol* 148:818–828
- Maciejewska B, Kopcewicz J (2002) Inhibitory effect of methyl jasmonate on flowering and elongation growth in *Pharbitis nil*. *J Plant Growth Regul* 21:216–223
- Maciejewska BD, Kęsy J, Zielińska M, Kopcewicz J (2004) Jasmonates inhibit flowering in short-day plant *Pharbitis nil*. *Plant Growth Regul* 43:1–8
- Martin JL, McMillan FM (2002) SAM (dependent) I AM: the S-adenosylmethionine-dependent methyltransferase fold. *Curr Opin Struct Biol* 12:783–793
- McKersie BD, Lepock JR, Kruuv J, Thompson JE (1987) The effects of cotyledon senescence on the composition and physical properties of membrane lipid. *Biochim Biophys Acta* 508:197–212
- Neill S, Barros R, Bright J, Desikan R, Hancock J, Harrison J, Morris P, Ribeiro D, Wilson I (2008) Nitric oxide, stomatal closure, and abiotic stress. *J Exp Bot* 59:165–176
- Noir S, Bömer M, Takahashi N, Ishida T, Tsui TL, Balbi V, Shanahan H, Sugimoto K, Devoto A (2013) Jasmonate controls leaf growth by repressing cell proliferation and the onset of

- endoreduplication while maintaining a potential stand-by mode. *Plant Physiol* 161:1930–1951
- Norastehnia A, Sajedi RH, Nojavani-Asghari M (2007) Inhibitory effects of methyl jasmonate on seed germination in maize (*Zea mays*): effect on α -amylase activity and ethylene production. *Gen Appl Plant Physiol* 33:13–23
- Oh SA, Lee SY, Chung IK, Lee CH, Nam HG (1996) A senescence-associated gene of *Arabidopsis thaliana* is distinctively regulated during natural and artificially induced leaf senescence. *Plant Mol Biol* 30:739–754
- Okushima Y, Mitina I, Quach HL, Theologis A (2005) AUXIN RESPONSE FACTOR 2 (ARF2): a pleiotropic developmental regulator. *Plant J* 43:29–46
- Pak H, Guo Y, Chen M, Chen K, Li Y, Hua S, Shamsi I, Meng H, Shi C, Jiang L (2009) The effect of exogenous methyl jasmonate on the flowering time, floral organ morphology, and transcript levels of a group of genes implicated in the development of oilseed rape flowers (*Brassica napus* L.). *Planta* 231:79–91
- Peterman TK, Siedow JN (1985) Behavior of lipoxygenase during establishment, senescence, and rejuvenation of soybean cotyledons. *Plant Physiol* 78:690–695
- Pružinská A, Tanner G, Aubry S, Anders I, Moser S, Müller T, Ongania KH, Kräutler B, Youn JY, Liljegren SJ, Hörtensteiner S (2005) Chlorophyll breakdown in senescent *Arabidopsis* leaves. Characterization of chlorophyll catabolites and of chlorophyll catabolic enzymes involved in the degreening reaction. *Plant Physiol* 139:52–56
- Quirino BF, Noh YS, Himelblau E, Amasino RM (2000) Molecular aspects of leaf senescence. *Trends Plant Sci* 5:278–282
- Reinbothe S, Reinbothe C, Parthier B (1993) Methyl jasmonate represses translation initiation of a specific set of mRNAs in barley. *Plant J* 4:459–467
- Reinbothe C, Parthier B, Reinbothe S (1997) Temporal pattern of jasmonate-induced alterations in gene expression of barley leaves. *Planta* 201:281–287
- Rukes KL, Mulkey TJ (1993) Hormonal regulation of cotyledon senescence. *Bioscience* 19:10–15
- Seo HS, Song JT, Cheong JJ, Lee YH, Lee YW, Hwang I, Lee JS, Choi YD (2001) Jasmonic acid carboxyl methyltransferase: a key enzyme for jasmonate-regulated plant responses. *Proc Natl Acad Sci USA* 98:4788–4793
- Staswick P (1992) Jasmonate, genes and fragrant signals. *Plant Physiol* 99:804–807
- Troncoso-Poncea MA, Caoc X, Yangb Z, Ohlrogge JB (2013) Lipid turnover during senescence. *Plant Sci* 205–206:13–19
- Tsai FY, Hung KT, Kao CH (1996) An increase in ethylene sensitivity is associated with jasmonate-promoted senescence of detached rice leaves. *J Plant Growth Regul* 15:197–200
- Ueda J, Kato J (1981) Promotive effect of methyl jasmonate on oat leaf senescence in the light. *Z Pflanzenphysiol* 103:357–359
- Van der Graaff E, Schwacke R, Schneider A, Desimone M, Flugge UI, Kunze R (2006) Transcription analysis of *Arabidopsis* membrane transporters and hormone pathways during developmental and induced leaf senescence. *Plant Physiol* 141:776–792
- Wada S, Ishida H, Izumi M, Yoshimoto K, Ohsumi Y, Mae T, Makino A (2009) Autophagy plays a role in chloroplast degradation during senescence in individually darkened leaves. *Plant Physiol* 149:885–893
- Wang CY (1998) Methyl jasmonate inhibits postharvest sprouting and improves storage quality of radishes. *Postharvest Biol Technol* 14:179–183
- Wasternack C, Hause B (2002) Jasmonates and octadecanoids: signals in plant stress responses and plant development. *Prog Nucleic Acid Res Mol Biol* 72:165–221
- Wasternack C, Kombrink E (2010) Jasmonates: structural requirements for lipid-derived signals active in plant stress responses and development. *ACS Chem Biol* 5:63–77
- Watanabe A, Hamada K, Yokoi H, Watanabe A (1994) Biphasic and differential expression of cytosolic glutamine synthetase genes of radish during seed germination and senescence of cotyledons. *Plant Mol Biol* 26:1807–1817
- Weaver LM, Gan S, Quirino B, Amasino RM (1998) A comparison of the expression patterns of several senescence-associated genes in response to stress and hormone treatment. *Plant Mol Biol* 37:455–469
- Weidhase RA, Kramell HM, Lehmann J, Liebisch HW, Lerbs W, Parthier B (1987a) Methyl jasmonate-induced changes in the polypeptide pattern of senescing barley leaf segments. *Plant Sci* 51:177–186
- Weidhase RA, Lehmann J, Kramell H, Sembdner G, Parthier B (1987b) Degradation of ribulose-15-bisphosphate-carboxylase and chlorophyll in senescing barley leaf segments triggered by jasmonic acid methyl ester and counteraction by cytokinin. *Physiol Plant* 69:161–166
- Weng JH (2006) Underestimate of PS2 efficiency in the field due to high leaf temperature resulting from leaf clipping and its amendment. *Photosynthetica* 44:467–470
- Wilmowicz E, Kęsy J, Kopcewicz J (2008) Ethylene and ABA interactions in the regulation of flower induction in *Pharbitis nil*. *J Plant Physiol* 165:1917–1928
- Wilmowicz E, Frankowski K, Grzegorzewska W, Kęsy J, Kućko A, Banach M, Szmidt-Jaworska A, Saniewski M (2014) The role of jasmonates in the formation of a compound of chalcones and flavans with phytoalexin-like properties in mechanically wounded scales of *Hippeastrum* × hybr. bulbs. *Acta Biol Crac Ser Bot* 56:54–58
- Wilson ZA, Song J, Taylor B, Yang C (2011) The final split: the regulation of anther dehiscence. *J Exp Bot* 62:1633–1649
- Woo HR, Chung KM, Park JH, Oh SA, Ahn T, Hong SH, Jang SK, Nam HG (2001) ORE9, an F-box protein that regulates leaf senescence in *Arabidopsis*. *Plant Cell* 13:1779–1790
- Wrisher M, Prebeg T, Magnus V, Ljubešić N (2009) Unusual thylakoid structures appearing during degradation of the photosynthetic apparatus in chloroplasts. *Acta Bot Croat* 68:1–9
- Xuan Y, Zhou S, Wang L, Cheng Y, Zhao L (2010) Nitric oxide functions as a signal and acts upstream of AtCaM3 in thermotolerance in *Arabidopsis* seedlings. *Plant Physiol* 153:1895–1906
- Zhang MP, Zhang CJ, Yu GH, Jiang YZ, Strasser RJ, Yuan ZY, Yang XS, Chen GX (2010) Changes in chloroplast ultrastructure, fatty acid components of thylakoid membrane and chlorophyll *a* fluorescence transient in flag leaves of a super-high-yield hybrid rice and its parents during the reproductive stage. *J Plant Physiol* 167:277–285
- Zhao MG, Chen L, Zhang LL, Zhang WH (2009) Nitric reductase-dependent nitric oxide production is involved in cold acclimation and freezing tolerance in *Arabidopsis*. *Plant Physiol* 151:755–767